

Avoidable Mortality

2020

Published on 7 March 2022



This report analyses deaths that were registered in Scotland in 2020 which are classified as avoidable by the Organisation for Economic Co-operation and Development (OECD).

It also includes breakdowns by sex, age, cause of death and deprivation.

The rate of avoidable mortality rose by 9% in 2020

In 2020 the age-standardised rate of avoidable mortality rose to 336 per 100,000 people (up by 9%). COVID-19 deaths aged under 75 helped to drive this increase.

Avoidable mortality rate (deaths per 100,000)




Avoidable mortality is higher in more deprived areas

In the most deprived 20% of areas in Scotland, the rate of avoidable mortality was 3.9 times as large as in the 20% least deprived areas.

For all causes of death, the rate of mortality is 1.9 times as high in the most deprived areas.

Avoidable mortality rate by SIMD quintile



Cancers and circulatory diseases had the highest rates of avoidable mortality

The rate of avoidable mortality for cancer deaths in 2020 was 98 per 100,000 people. The rate for preventable cancer deaths was more than twice that of treatable cancer deaths.

Avoidable mortality rate (deaths per 100,000), 2020




Table of Contents

1. Key Findings	4
2. Introduction.....	5
3. Avoidable mortality in Scotland	6
4. Avoidable mortality by deprivation	8
5. Avoidable mortality by cause	10
6. Avoidable Mortality in Great Britain.....	13
7. Avoidable mortality by Health Board	14
8. Methodology	15
9. Strengths and Limitations	15
10. Future developments	16
11. Related Statistics.....	16
12. Notes on statistical publications.....	16

1. Key Findings

- There were 64,093 registered deaths in Scotland in 2020, of which 17,153 (27%) were considered avoidable.
- The rate of avoidable deaths in 2020 was 9% higher than the previous year. This increase was largely due to COVID-19 deaths.
- After adjusting for age, avoidable mortality rates among males (425 per 100,000) were 1.7 times as high as among females (253 per 100,000).
- Cancers and circulatory diseases were the most common causes of avoidable mortality in 2020, accounting for 29% and 25% of all avoidable deaths respectively.
- Alcohol and drug related avoidable mortality rates increased for the ninth year in a row, with 52 deaths per 100,000 people.
- After adjusting for age, avoidable mortality rates in the most deprived areas were nearly 4 times the rates of those in the least deprived areas.
- Scotland has a higher avoidable mortality rate than both England and Wales, as well as the GB average. This continues the historic trend of Scotland having the highest avoidable mortality rates among GB countries.

2. Introduction

What is Avoidable Mortality?

Deaths which are considered either preventable or treatable through public health interventions or timely and effective healthcare.

It is based on an [international definition by the OECD/Eurostat](#) and looks only at deaths under the age of 75.

When discussing avoidable deaths, the following terms are used:

Preventable mortality – deaths that can be mainly avoided through effective public health and primary prevention interventions

Treatable mortality – deaths that can be mainly avoided through timely and effective healthcare interventions, including secondary prevention and treatment

Avoidable mortality – deaths defined as either preventable or treatable

Why use age-standardised mortality rates?

Age-standardised mortality rates are a better measure of mortality than numbers of deaths, as they account for the population size and age structure and provide more reliable comparisons between groups or over time.

More information on the calculation of age-standardised mortality rates is available on our [website](#).


3. Avoidable mortality in Scotland

The rate of avoidable deaths in 2020 was 9% higher than the previous year. This increase was largely due to COVID-19 deaths.

Of the 64,093 deaths registered in 2020:

- 12,294 (19% of all registered deaths) were considered preventable.
- 4,860 (8% of all registered deaths) were as a result of a treatable cause.
- 17,153 (27% of all registered deaths) in total were considered avoidable¹.

Figure 1: Avoidable mortality rates in Scotland: 2001-2020



The rate of avoidable mortality increased by 9% in 2020. This is the first statistically significant increase since 2015 when the rate increased by 3% (Figure 1).


The rates for both males and females increased in 2020, with the rate for males remaining higher (1.7 times as high in 2020) than for females.

A large jump in preventable deaths occurred in 2020, and the number of male preventable deaths remains higher than female preventable deaths. The difference between male and female treatable deaths is smaller (Figure 2).

¹ Figures for preventable and treatable causes of deaths may not sum to the total for avoidable deaths due to rounding.

Figure 1 and Figure 2 show figures without 2020 COVID-19 deaths (dotted line). It illustrates that the increase in overall avoidable mortality in 2020 was largely due to the inclusion of COVID-19 as a preventable death by the OECD.

Figure 2: Preventable and treatable deaths, by sex: 2001-2020



4. Avoidable mortality by deprivation

People living in the most deprived areas in Scotland are 3.9 times as likely to die an avoidable death compared to those in the least deprived areas. The increase in avoidable mortality rates in 2020 has disproportionately occurred in the most deprived areas.

Figure 3 shows avoidable mortality rates in each SIMD (Scottish index of multiple deprivation) quintiles over the last two decades.

SIMD

The Scottish index multiple deprivation is a measure of how deprived an area is. A score is given to all of Scotland's datazones based on multiple indicators of deprivation.


The datazones are then ranked 1 to 6,976 based on their score. Subsequently, the rankings are split into 5 equally sized groups forming SIMD quintiles.

All quintiles experienced an increase in avoidable mortality in 2020. However, areas of greater deprivation experienced a larger effect.

In 2020, the avoidable mortality rate of those in quintiles 1 (the 20% most deprived areas) was 634 per 100,000 people compared with 162 per 100,000 for those in quintile 5 (the 20% least deprived areas).

The most deprived quintile was 3.9 times as likely to die from an avoidable death in 2020 compared to those in the least deprived quintile. The increase in avoidable mortality seen in 2020 disproportionately affected the most deprived.

Figure 3: Avoidable mortality rates by deprivation, all persons: 2001-2020



5. Avoidable mortality by cause

Cancers and circulatory system diseases were the leading causes of avoidable mortality in 2020, while alcohol and drug related deaths continue to increase.

The [OECD/Eurostat definition](#) includes COVID-19 (see new category “Provisional assignment of new diseases”) added as a preventable mortality.

The leading causes of avoidable deaths in 2020 for all persons (see Figure 4) were:

- Cancers 5,043 avoidable deaths (a rate of 98 per 100,000 people).
- Circulatory system diseases (4,270 avoidable deaths, a rate of 83 per 100,000 people).
- Alcohol and drug related disorders (2,576 avoidable deaths, a rate of 52 per 100,000 people).

Not all causes of death are avoidable

The proportion of deaths that are classified as avoidable varies greatly between ICD-10 chapters. For example, almost all deaths due to external causes are counted as avoidable, whereas under one tenth of deaths due to mental disorders hold this classification. See **Table A** for details.

For males the leading cause of avoidable and treatable deaths were circulatory system diseases, whilst the leading cause of preventable deaths were cancers.

In comparison, for females the leading cause of all avoidable, preventable and treatable deaths were cancers.

COVID-19 was a significant contributor to avoidable mortality in 2020, with an avoidable mortality rate of 29 per 100,000 people. Males had a higher rate of avoidable deaths, with 37 deaths per 100,000 versus females with 21 deaths per 100,000.

Figure 4: Avoidable mortality rates in Scotland, by cause: 2020





Figure 5: Avoidable mortality rate: select causes 2001-2020



Avoidable mortality rates from cancers (the largest contributor) have remained relatively stable over the last 20 years (see figure 5). The next largest contributor,

Circulatory system diseases, saw a slight increase in 2020 after remaining stable for six years.

Alcohol and drug related disorders increased in 2020, furthering a trend since 2012 of increasing avoidable mortality.


For a full summary of all causes of death in Scotland (not only avoidable mortality) please see [Section 6: Deaths - Causes in the NRS Vital Events Reference Tables 2020](#).

6. Avoidable Mortality in Great Britain

Scotland continues to have higher avoidable mortality than both England and Wales.

Figure 6 compares the avoidable mortality rates between Scotland, England, Wales, and the average for Great Britain (GB)¹. Scotland (336 deaths per 100,000 people) has a higher avoidable mortality rate than both England (257 deaths per 100,000 people) and Wales (287 deaths per 100,000 people), as well as the GB average (266 deaths per 100,000 people). This continues the historic trend of Scotland having higher avoidable mortality rates, although the gap between Scotland and the GB average narrowed slightly in the 2020. All GB nations experienced statistically significant increases in 2020.

Figure 6: Avoidable mortality rates in Great Britain (GB): 2001-2020




¹ GB data from the ONS. Deaths of non-residents are excluded for England and Wales and included for GB and Scotland. These statistics are also based on the OECD definition of avoidable mortality. See the [ONS Avoidable Mortality bulletin](#) and [dataset](#) for further detail. UK and Northern Ireland (NI) data will be released at a later date pending the outcome of the current review of suicide statistics in NI, as these deaths form part of the definition of avoidable mortality (see ONS bulletin for further detail).

7. Avoidable mortality by Health Board

Health boards in the west of Scotland experienced the highest rates of avoidable mortality.

Figure 7: Avoidable mortality rates by health board, all persons: 2018-2020 average



The highest avoidable mortality rates were recorded in Greater Glasgow and Clyde (388 deaths per 100,000), Lanarkshire (359 deaths per 100,000), and in Ayrshire and Arran (350 deaths per 100,000).

Shetland (222 deaths per 100,000) and Orkney (230 deaths per 100,000) saw the lowest avoidable mortality rates.

8. Methodology

NRS has updated its definition of avoidable mortality to use the [new OECD international definition of avoidable mortality](#) to ensure our statistics are comparable internationally and with other parts of the UK. The new definition has been applied to the full time series back to 2001.

Deaths are classified as preventable or treatable according to their cause of death and the age at death.

9. Strengths and Limitations

Strengths

1. Information is supplied when a death is registered, giving complete coverage of the population and ensures highly accurate estimates that are representative of the population.
2. Coding for cause of death is carried out according to the World Health Organization (WHO) ICD-10 framework.
3. The use of age-standardised mortality rates means our statistics are comparable between local health boards, councils and at national level.
4. The use of the OECD definition also makes the statistics in this report comparable internationally.

Limitations

1. Data are insufficiently robust to provide local authority estimates for smaller areas for single years (hence why they are aggregated over a 3-year period). This limits the timeliness of the data.
2. In a small number of cause of death categories, the number of deaths was too small to report a reliable rate. For this reason, the categories were combined in and thus limits the precision accuracy of our cause of death reporting.

10. Future developments

The next scheduled publication for further avoidable mortality statistics in Scotland is due late 2022.

11. Related Statistics

- Prior to 2019, avoidable mortality statistics (using a previous definition) were published in a summary format on the [NRS website](#).
- The [ONS](#) publish avoidable mortality statistics on the UK and its constituent countries (including data included in this report in section 6). These are also based on the OECD definition of avoidable mortality. See the [ONS Avoidable Mortality bulletin](#) and [dataset](#) for further detail.

12. Notes on statistical publications

National Statistics

The United Kingdom Statistics Authority (UKSA) has designated these statistics as National Statistics, in line with the Statistics and Registration Service Act 2007 and signifying compliance with the Code of Practice for Official Statistics (available on the [UKSA](#) website).

National Statistics status means that official statistics meet the highest standards of trustworthiness, quality and public value.

All official statistics should comply with all aspects of the Code of Practice for Official Statistics. They are awarded National Statistics status following an assessment by the Authority's regulatory arm. The Authority considers whether the statistics meet the highest standards of Code compliance, including the value they add to public decisions and debate.

It is National Records of Scotland's responsibility to maintain compliance with the standards expected of National Statistics. If we become concerned about whether these statistics are still meeting the appropriate standards, we will discuss any concerns with the Authority promptly. National Statistics status can be removed at any point when the highest standards are not maintained, and reinstated when standards are restored.

Information on background and source data

Further details on data source(s), timeframe of data and timeliness, continuity of data, accuracy, etc. can be found in the metadata that is published alongside this publication on the NRS website.

National Records of Scotland

We, the National Records of Scotland, are a non-ministerial department of the devolved Scottish Administration. Our aim is to provide relevant and reliable information, analysis and advice that meets the needs of government, business and the people of Scotland. We do this as follows:

Preserving the past – We look after Scotland’s national archives so that they are available for current and future generations, and we make available important information for family history.

Recording the present – At our network of local offices, we register births, marriages, civil partnerships, deaths, divorces and adoptions in Scotland.

Informing the future – We are responsible for the Census of Population in Scotland which we use, with other sources of information, to produce statistics on the population and households.

You can get other detailed statistics that we have produced from the [Statistics](#) section of our website. Scottish Census statistics are available on the [Scotland’s Census](#) website.

We also provide information about [future publications](#) on our website. If you would like us to tell you about future statistical publications, you can register your interest on the Scottish Government [ScotStat website](#).

You can also follow us on twitter [@NatRecordsScot](#)

Enquiries and suggestions

Please get in touch if you need any further information, or have any suggestions for improvement.

E-mail: statisticscustomerservices@nrscotland.gov.uk

Lead Statistician: Julie Ramsay