

Table 1**Increased winter mortality^{1,2} by age group, Scotland, 1990/1991 to 2007/2008**

	0-64	65-74	75-84	85+	All ages
1990/91	230	580	750	880	2,430
1991/92	350	560	1,020	950	2,890
1992/93	280	550	950	960	2,740
1993/94	350	440	990	800	2,590
1994/95	240	380	930	760	2,310
1995/96	250	860	1,420	1,120	3,650
1996/97	320	630	1,350	1,350	3,640
1997/98	170	730	950	760	2,610
1998/99	380	790	1,660	1,920	4,750
1999/2000	650	970	1,820	1,750	5,190
2000/01	260	370	820	760	2,220
2001/02	80	230	820	710	1,840
2002/03	350	300	940	920	2,510
2003/04	320	510	840	1,170	2,840
2004/05	200	430	1,030	1,090	2,760
2005/06	330	280	550	610	1,780
2006/07	190	410	980	1,180	2,750
2007/08 prov.	130	320	880	850	2,180

Notes

1. Increased winter mortality has been defined as the difference between the number of deaths in the months December - March and the average of the preceding (August - November) and following (April - July) non-winter periods.

2. Because of the approximate nature of this measure, numbers have been rounded independently to the nearest 10. The sum of the age group figures may, therefore, differ from the "all ages" total.