

Table 4.7 Infant death rates, by sex and cause, Scotland, 1995 to 2005

| ICD Codes | | Cause of death | Infant deaths per 1,000 live births | | | | | | | | | | | | |
|-----------------|----------------|---|-------------------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| ICD9 list | ICD10 list | | 1995 | 1996 | 1997 | 1998 | 1999 | | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | |
| | | | | | | | ICD9 | ICD10 | | | | | | | |
| | | All causes | P | 6.2 | 6.2 | 5.3 | 5.6 | 5.0 | 5.0 | 5.7 | 5.5 | 5.3 | 5.1 | 4.9 | 5.2 |
| | | | M | 6.4 | 6.7 | 6.1 | 6.2 | 5.7 | 5.7 | 6.4 | 5.8 | 6.4 | 5.4 | 5.8 | 5.7 |
| | | | F | 6.1 | 5.5 | 4.5 | 4.9 | 4.3 | 4.3 | 5.1 | 5.2 | 4.1 | 4.7 | 4.0 | 4.8 |
| 001-739 | A00-Q99 | All causes in Chapters I - XV | M | 0.7 | 0.5 | 0.6 | 0.8 | 0.4 | 0.5 | 0.4 | 0.4 | 0.6 | 0.3 | 0.5 | 0.5 |
| | | | F | 0.6 | 0.6 | 0.6 | 0.4 | 0.5 | 0.9 | 0.6 | 0.2 | 0.3 | 0.3 | 0.6 | 0.5 |
| 760-779 | P00-P96 | XVI. Certain conditions originating in the perinatal period | M | 2.9 | 3.4 | 2.7 | 3.3 | 2.9 | 2.8 | 3.5 | 3.3 | 3.4 | 3.0 | 3.3 | 3.5 |
| | | - of which | F | 3.0 | 2.6 | 2.0 | 2.4 | 2.0 | 1.7 | 2.8 | 3.0 | 2.6 | 2.6 | 2.2 | 2.5 |
| 764-766 | P05-08 | Disorders related to length of gestation and fetal growth | M | 0.4 | 0.5 | 0.4 | 0.6 | 0.5 | 0.4 | 0.8 | 0.6 | 0.3 | 0.7 | 0.8 | 0.5 |
| | | | F | 0.4 | 0.5 | 0.4 | 0.5 | 0.4 | 0.4 | 0.5 | 0.7 | 0.3 | 0.6 | 0.3 | 0.4 |
| 768-770 | P20-29 | Respiratory and cardiovascular disorders specific to the perinatal period | M | 1.5 | 1.7 | 1.3 | 1.7 | 1.2 | 1.4 | 1.1 | 1.1 | 1.0 | 0.5 | 1.0 | 1.0 |
| | | | F | 1.3 | 1.3 | 1.0 | 1.0 | 0.6 | 0.7 | 1.0 | 0.4 | 0.7 | 0.7 | 0.7 | 0.5 |
| 740-759 | Q00-Q99 | XVII. Congenital malformations deformations and chromosomal abnormalities | M | 1.8 | 1.9 | 1.5 | 1.2 | 1.5 | 1.4 | 1.5 | 1.4 | 1.3 | 1.0 | 1.2 | 1.4 |
| | | | F | 1.6 | 1.7 | 1.3 | 1.3 | 1.0 | 0.9 | 1.0 | 1.3 | 0.8 | 1.1 | 0.8 | 1.2 |
| 780-799 | R00-R99 | XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified | M | 0.8 | 1.0 | 1.1 | 0.6 | 0.8 | 0.8 | 0.7 | 0.6 | 0.9 | 1.1 | 0.6 | 0.3 |
| | | - of which | F | 0.7 | 0.5 | 0.6 | 0.6 | 0.7 | 0.7 | 0.5 | 0.7 | 0.3 | 0.6 | 0.4 | 0.5 |
| 798.0 | R95 | Sudden infant death syndrome | M | 0.8 | 1.0 | 1.1 | 0.6 | 0.8 | 0.8 | 0.7 | 0.6 | 0.9 | 1.1 | 0.6 | 0.3 |
| | | | F | 0.7 | 0.5 | 0.6 | 0.6 | 0.7 | 0.7 | 0.5 | 0.7 | 0.3 | 0.5 | 0.4 | 0.5 |
| E800-999 | V01-Y98 | XX. External causes of morbidity and mortality | M | 0.2 | 0.0 | 0.2 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | - | 0.1 | 0.1 |
| | | | F | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 | - | 0.1 |